

GEOMETRY OF THE AHARONOV-BOHM EFFECT

R. S. Huerfano^{1,2}, M. A. López² and M. Socolovsky²

¹ *Departamento de Matemáticas, Universidad Nacional de Colombia, Bogotá, Colombia*

² *Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México
Circuito Exterior, Ciudad Universitaria, 04510, México D. F., México*

We show that the connection responsible for any abelian or non abelian Aharonov-Bohm effect with n parallel “magnetic” flux lines in $\mathbb{R}^3$, lies in a trivial G -principal bundle $P \rightarrow M$, i.e. P is isomorphic to the product $M \times G$, where G is any path connected topological group; in particular a connected Lie group. We also show that two other bundles are involved: the universal covering space $\tilde{M} \rightarrow M$, where path integrals are computed, and the associated bundle $P \times_G \mathbb{C}^m \rightarrow M$, where the wave function and its covariant derivative are sections.

Key words: Aharonov-Bohm effect; fibre bundle theory; gauge invariance.

As is well known, the magnetic Aharonov-Bohm (A-B) effect ^{1,2} is a gauge invariant, non local quantum phenomenon, with gauge group $U(1)$, which takes place in a non simply connected space. It involves a magnetic field in a region where an electrically charged particle obeying the Schroedinger equation cannot enter, i.e. the ordinary 3-dimensional space minus the space occupied by the solenoid producing the field; in the ideal mathematical limit, the solenoid is replaced by a flux line. Locally, the particle couples to the magnetic potential $\vec{A}$ but not to the magnetic field $\vec{B}$; however, the effect is gauge invariant since it only depends on the flux of $\vec{B}$ inside the solenoid.

The fibre bundle theoretic description of this kind of phenomena has proved to be very useful to obtain a more profound insight into the relation between physical processes and pure mathematics. ³ In the present case, since by symmetry, the dimension along the flux line can be ignored, the problem reduces to the effect on the charged particle of an abelian connection in a $U(1)$ -bundle with base space the plane minus a point. The wave function representing the particle is a section of an associated vector bundle. As shown in ref. 4, the bundle turns out to be trivial and then its total space is isomorphic to the product $(\mathbb{R}^2 - \{point\}) \times U(1)$. Since $\mathbb{R}^2$ is topologically equivalent to an open disk, and $U(1)$ is the unit circle, the bundle structure is summarized by

$$U(1) \rightarrow T_{o*}^2 \rightarrow D_{o*}^2 \quad (1)$$

where D_{o*}^2 is the open disk minus a point and T_{o*}^2 is the open solid 2-torus minus a circle.

As suggested by Wu and Yang ⁶, Yang-Mills fields can give rise to non abelian A-B effects. In ref. 6 the authors studied an $SU(2)$ gauge configuration leading to an A-B effect; later, several authors studied the effect with gauge groups $SU(3)$ ⁷ and $U(N)$ ⁸. Also, in refs. 9-14 the effect was studied in the context of gravitation theory. In all these examples, as in the magnetic case, there is a principal bundle structure

$$\xi : G \rightarrow P \xrightarrow{\pi_G} M \quad (2)$$

whose total space P , where the connection giving rise to the effect lies, is however never specified. We shall restrict ourselves to connections ω giving rise to n ($n = 1, 2, 3, \dots$) flux lines in $\mathbb{R}^3$. In this note we prove the following:

Theorem: Let G be a path connected topological group (for example a connected Lie group) and ξ a continuous principal G -bundle over $\mathbb{R}^3 - \{n \text{ parallel lines}\}$, $n = 1, 2, 3, \dots$. Then the bundle ξ is trivial, i.e. isomorphic to the product bundle.

Proof of the theorem

The classification of bundles over $\mathbb{R}^3 \setminus \{n \text{ parallel lines}\}$ is the same as that over $\mathbb{R}^2 \setminus \{n \text{ points}\}$ which is topologically equivalent to $D_{\circ}^2 \setminus \{n \text{ points}\} \equiv D_{\circ * n}^2$. Denote this set of points by $\{b_1, \dots, b_n\}$. By symmetry along the dimension of the flux tubes, $D_{\circ * n}^2$ is the space where it can be considered that the charged particles move.

Let x_0 be a point in $D_{\circ * n}^2$. We construct a bouquet of n loops $\gamma_1, \gamma_2, \dots, \gamma_n$ through x_0 , with the k -th loop surrounding the point b_k , $k = 1, 2, \dots, n$. This space is homeomorphic to the wedge product (or reduced join) $S^1 \vee \dots \vee S^1 \equiv \vee_n S^1 \equiv S_{(1)}^1 \vee \dots \vee S_{(n)}^1$ of n circles¹⁵, and the classification of bundles over $D_{\circ * n}^2$ is the same as that over $\vee_n S^1$, namely

$$\mathcal{B}_{D_{\circ * n}^2}(G) = \mathcal{B}_{\vee_n S^1}(G) \quad (3)$$

where $\mathcal{B}_M(G)$ is the set of isomorphism classes of G -bundles over M ¹⁶.

By explicit construction we shall prove that, up to isomorphism, the unique G -bundle over $\vee_n S^1$ is the product bundle $\vee_n S^1 \times G$ (which is a purely topological result). With this aim, we cover the circle $S_{(k)}^1$ with two open sets U_{k+} and U_{k-} such that

$$U_{k+} \cap U_{k-} \simeq \{x_0, a_k\}, \quad k = 1, \dots, n, \quad (4)$$

$$U_{i+} \cap U_{j+} \simeq U_{i-} \cap U_{j-} \simeq U_{i+} \cap U_{j-} \simeq \{x_0\}, \quad i, j = 1, \dots, n, \quad i \neq j \quad (5)$$

where $\simeq$ denotes homotopy equivalence, and $a_k \in S_{(k)}^1$ with $a_k \neq x_0$. We then have

$$2 \times \binom{2n}{2} + 2n = \frac{2 \times (2n)!}{(2n-2)!2!} + 2n = 4n^2 \quad (6)$$

transition functions

$$g_{\alpha, \beta} : U_{\alpha} \cap U_{\beta} \rightarrow G \quad (7)$$

with $g_{\beta, \alpha} = g_{\alpha, \beta}^{-1}$. Up to homotopy they are given by

$$g_{k+, k-} : \{x_0, a_k\} \rightarrow G, \quad x_0 \mapsto g_{0k}, \quad a_k \mapsto g_k, \quad k = 1, \dots, n \quad (8)$$

and

$$\begin{aligned} g_{i+, j+}, g_{i-, j-}, g_{i+, j-} : \{x_0\} &\rightarrow G, \quad i, j = 1, \dots, n, \quad i \neq j, \\ x_0 &\mapsto g_{ij++}, \quad x_0 \mapsto g_{ij--}, \quad x_0 \mapsto g_{ij+-}. \end{aligned} \quad (9)$$

The $2n$ transition functions $g_{i+, i+}, g_{i-, i-}$, $i \in \{1, \dots, n\}$, give the identity in G . The number of cocycle relations $g_{\beta, \alpha} g_{\alpha, \gamma} = g_{\beta, \gamma}$ on the $2n(2n-1)$ non trivial transition functions $g_{\mu, \nu}$ is $\binom{2n}{3} = \frac{n(2n-1)(2n-2)}{3}$.

Let $g'_{k+, k-} : \{x_0, a_k\} \rightarrow G$ be given by $g'_{k+, k-}(x_0) = g'_{0k}$, $g'_{k+, k-}(a_k) = g'_k$. Since G is path connected, there exist continuous paths

$$c_0^{(k)} : [0, 1] \rightarrow G, \quad t \mapsto c_0^{(k)}(t) \quad \text{with} \quad c_0^{(k)}(0) = g_{0k}, \quad c_0^{(k)}(1) = g'_{0k}$$

and

$$c_k^{(k)} : [0, 1] \rightarrow G, \quad t \mapsto c_k^{(k)}(t) \quad \text{with} \quad c_k^{(k)}(0) = g_k, \quad c_k^{(k)}(1) = g'_k.$$

Then the continuous function

$$H_k : \{x_0, a_k\} \times [0, 1] \rightarrow G$$

given by

$$H_k(x_0, t) = c_0^{(k)}(t), \quad H_k(a_k, t) = c_k^{(k)}(t) \quad (10)$$

is a homotopy between $g_{k+, k-}$ and $g'_{k+, k-}$ since

$$H_k(x_0, 0) = g_{0k} \quad \text{and} \quad H_k(a_k, 0) = g_k$$

i.e.

$$H_k|_{\{x_0, a_k\} \times \{0\}} = g_{k+, k-}$$

and

$$H_k(x_0, 1) = g'_{0k} \quad \text{and} \quad H_k(a_k, 1) = g'_k$$

i.e.

$$H_k|_{\{x_0, a_k\} \times \{1\}} = g'_{k+, k-}.$$

Then the homotopy class of maps from $\{x_0, a_k\}$ to G has only one element, namely $[g_{k+, k-}] \sim$ i.e.

$$[\{x_0, a_k\}, G] \sim = \{[g_{k+, k-}] \sim\}. \quad (11)$$

Similarly, let $g_{i\mu, j\nu}$ with $\mu, \nu \in \{+, -\}$ be any of the functions in (9), and let $g'_{i\mu, j\nu} : \{x_0\} \rightarrow G$ be given by

$$g'_{i\mu, j\nu}(x_0) = g'_{ij, \mu\nu}; \quad (12)$$

then the map

$$H_{i\mu, j\nu} : \{x_0\} \times [0, 1] \rightarrow G, \quad H_{i\mu, j\nu}(x_0, t) = c_{i\mu, j\nu}(t) \quad (13)$$

with $c_{i\mu, j\nu} : [0, 1] \rightarrow G$ a continuous path in G satisfying $c_{i\mu, j\nu}(0) = g_{ij\mu\nu}$ and $c_{i\mu, j\nu}(1) = g'_{ij\mu\nu}$, is a homotopy between $g_{i\mu, j\nu}$ and $g'_{i\mu, j\nu}$ i.e.

$$[\{x_0\}, G] \sim = \{[g_{i\mu, j\nu}] \sim\}. \quad (14)$$

It is then easy to show that, if we define (constant) functions

$$\Lambda_{k\mu} : U_{k\mu} \rightarrow G, \quad p \mapsto \lambda_{k\mu}, \quad k = 1, \dots, n, \quad \mu = +, -, \quad (15)$$

then

$$g'_{k+, k-}(x_0) = \lambda_{k-} g_{k+, k-}(x_0) \lambda_{k+}^{-1}, \quad (16)$$

$$g'_{k+, k-}(a_k) = \lambda_{k-} g_{k+, k-}(a_k) \lambda_{k+}^{-1}, \quad (17)$$

for $k = 1, \dots, n$, and

$$g'_{i\mu, j\nu}(x_0) = \lambda_{j\nu} g_{i\mu, j\nu}(x_0) \lambda_{i\mu}^{-1} \quad (18)$$

for $i, j \in \{1, \dots, n\}$, $i \neq j$, $\mu, \nu \in \{+, -\}$. In fact, for arbitrary two sets of homotopic transition functions $g'_{\beta, \alpha} \sim g_{\beta, \alpha}$, one has continuous maps $H : (U_\beta \cap U_\alpha) \times [0, 1] \rightarrow G$ such that $H(x, 0) = g_{\beta, \alpha}(x)$ and $H(x, 1) = g'_{\beta, \alpha}(x)$. Given a map $\Lambda_\alpha : U_\beta \cap U_\alpha \rightarrow G$, $x \mapsto \Lambda(x) = \lambda_\alpha$, one defines $\bar{H}(x, 1) = H(x, 1)\Lambda_\alpha(x)$ and therefore $\bar{H}(x, 1) = g'_{\beta, \alpha}(x)\Lambda_\alpha(x)$. Now, one defines $\Lambda_\beta : U_\beta \cap U_\alpha \rightarrow G$ through $\Lambda_\beta(x) = \bar{H}(x, 1)H(x, 0)^{-1} = g'_{\beta, \alpha}(x)\Lambda_\alpha(x)g_{\beta, \alpha}(x)^{-1}$ i.e. $g'_{\beta, \alpha}(x) = \Lambda_\beta(x)g_{\beta, \alpha}(x)\Lambda_\alpha(x)^{-1}$ (*) for all $x \in U_\beta \cap U_\alpha$. In our case, for $U_{k+} \cap U_{k-} \simeq \{x_0, a_k\}$, the formulae corresponding to (*) are

$$g'_{k+, k-}(x_0) = \Lambda_{k-}(x_0)g_{k+, k-}(x_0)\Lambda_{k+}(x_0)^{-1}$$

and

$$g'_{k+, k-}(a_k) = \Lambda_{k-}(a_k)g_{k+, k-}(a_k)\Lambda_{k+}(a_k)^{-1}.$$

Extending continuously (as constants) Λ_{k+} and Λ_{k-} respectively to the open sets U_{k+} and U_{k-} :

$$\Lambda_{k+} : U_{k+} \rightarrow G, \quad p \mapsto \lambda_{k+}, \quad \Lambda_{k-} : U_{k-} \rightarrow G, \quad q \mapsto \lambda_{k-}$$

(in particular one has $\Lambda_{k+}(x_0) = \Lambda_{k+}(a_k) = \lambda_{k+}$ and $\Lambda_{k-}(x_0) = \Lambda_{k-}(a_k) = \lambda_{k-}$), one obtains equations (16) and (17). Proceeding similarly for the cases $U_{i\mu} \cap U_{j\nu} \simeq \{x_0\}$, one gets equation (18).

Then, up to isomorphisms, and according to a general theorem for coordinate bundles ¹⁷, there is a unique G -bundle generated by the transition functions given by the equations (8) and (9), namely the product bundle. We then have

$$\mathcal{B}_{\vee_n S^1}(G) = \{[G \rightarrow \vee_n S^1 \times G \rightarrow \vee_n S^1]\}, \quad (19)$$

where $[\]$ denotes here the equivalence class of bundles isomorphic to the product bundle. QED

For the cases of the examples in refs. 1, 6 and 7, we have, respectively,

$$\mathcal{B}_{D_{\circ*}^2}(U(1)) = \{[U(1) \rightarrow D_{\circ*}^2 \times U(1) \rightarrow D_{\circ*}^2]\}, \quad (20)$$

$$\mathcal{B}_{D_{\circ*}^2}(SU(2)) = \{[SU(2) \rightarrow D_{\circ*}^2 \times SU(2) \rightarrow D_{\circ*}^2]\}, \quad (21)$$

and

$$\mathcal{B}_{D_{\circ*2}^2}(SU(3)) = \{[SU(3) \rightarrow D_{\circ*2}^2 \times SU(3) \rightarrow D_{\circ*2}^2]\}. \quad (22)$$

In the gravitational case, the gauge group is $SL(2, \mathbb{C})$ ¹⁸, which is the universal covering group of the connected component of the Lorentz group $L_+^\uparrow$; then, for weak gravitational fields with a distribution of n gravitomagnetic flux lines as above we would have the trivial bundle

$$SL(2, \mathbb{C}) \rightarrow D_{\circ*n}^2 \times SL(2, \mathbb{C}) \rightarrow D_{\circ*n}^2. \quad (23)$$

We want to stress that, even if the A-B connection is flat (though not exact), one of the sufficient conditions for the automatic triviality of the A-B bundle fails: though paracompact, the bouquet $\vee_n S^1$ is not simply connected. (See corollary 9.2. in ref. 21, p. 92.) In addition, there is not a priori any physical reason why, under the specified conditions on the “magnetic flux”, the corresponding A-B bundle should be trivial.

It is interesting to notice that there are two other fibre bundles related to the A-B effect. The first bundle is the *universal covering space*¹⁹ of the base manifold (“laboratory” or physical space where the particles coupled to the A-B potential move) which is the $\pi_1(M; x_0)$ -(non trivial) bundle $\xi_c : \tilde{M} \xrightarrow{\pi} M$, where $\pi_1(M; x_0) (\equiv \pi_1(M)$ if M is connected) is the fundamental group of M . In our case,

$$\pi_1(\vee_n S^1; x_0) \cong \pi_1(\mathbb{R}^2 \setminus \{n \text{ points}\}; x_0) \cong \langle \{c_1, \dots, c_n\} \rangle$$

is the freely generated group with n non commuting generators $c_1, \dots, c_n$. In particular, for the original abelian A-B effect with group $U(1)$, $\mathbb{R}^{2*} = RS(Log)$: the Riemann surface of the logarithm, and $\pi_1(\mathbb{R}^{2*}) \cong \mathbb{Z}$.

The *particle propagator* in M , $K(x'', t''; x', t')$ with $t'' > t'$, is a sum of homotopy propagators⁷ multiplied by corresponding gauge factors²⁰: the former are given by unrestricted path integrals computed in $\tilde{M}$, the paths in these path integrals project onto the corresponding homotopy classes of paths in the non simply connected space M ; the latter are Wilson loops given by

$$Texp \int_{\pi(c)} \vec{A} \cdot d\vec{l}$$

where T denotes time order, $\vec{A}$ is the A-B potential, and c is a loop in $\tilde{M}$ beginning and ending respectively at y_0 and y'' in $\pi^{-1}(\{x''\}) \xrightarrow{\Psi} \pi_1(M)$, with y_0 fixed and arbitrary. Then one has the group homomorphism (many-to-one or one-to-one)

$$y'' \xrightarrow{\Psi} \Psi(y'') \xrightarrow{\varphi} Texp \int_{\pi(c)} \vec{A} \cdot d\vec{l} \in G \quad (24)$$

whose image in G , $\varphi(\pi_1(M))$, responsible for the A-B effect, is the *holonomy* of the connection.²¹

The second bundle is the *associated complex vector bundle* $\xi_{\mathbb{C}^m} : \mathbb{C}^m - P_{\mathbb{C}^m} \xrightarrow{\pi_{\mathbb{C}^m}} M$ ($m = 2s + 1$ is the dimension of the spinor space and s is the spin; for scalar particles $m = 1$), where

$$P_{\mathbb{C}^m} = (M \times G) \times_G \mathbb{C}^m = \{[(x, g), \vec{z}]\}_{((x, g), \vec{z}) \in (M \times G) \times_G \mathbb{C}^m},$$

$$[(x, g), \vec{z}] = \{(x, gg'), g'^{-1}\vec{z}\}_{g' \in G}. \quad (25)$$

$\xi_{\mathbb{C}^m}$ is trivial since ξ is trivial, and the quantum mechanical *wave functions* of the particles are global sections of $\xi_{\mathbb{C}^m}$:

$$\psi \in \Gamma(\xi_{\mathbb{C}^m})$$

i.e. $\psi : M \rightarrow P_{\mathbb{C}^m}$ with $\pi_{\mathbb{C}^m} \circ \psi = Id_M$.

Notice that while the propagator is computed in ξ_c , the wave function lies in $\xi_{\mathbb{C}^m}$, with

$$\psi(x'', t'') = \int_M dx' K(x'', t''; x', t') \psi(x', t'). \quad (26)$$

If ω is the A-B connection in P , then the coupling $\omega - \psi$ is the covariant derivative

$$\nabla_V^\omega \psi = \psi_{V^\uparrow(\gamma_\psi)} \in \Gamma(\xi_{\mathbb{C}^m}), \quad (27)$$

where V is a vector field in M , $V^\uparrow$ its horizontal lifting in P by ω , γ_ψ and $V^\uparrow(\gamma_\psi)$ are equivariant functions from P to $\mathbb{C}^m$ with $\gamma_\psi(p) = \vec{z}$ where $\psi(\pi_G(p)) = [p, \vec{z}]$, and $\psi_{V^\uparrow(\gamma_\psi)}(x) = [p, V^\uparrow(\gamma_\psi)(p)]$ for any $p \in \pi_G^{-1}(\{x\})$. Locally, of course, $\nabla_V^\omega \psi$ reproduces the usual minimal coupling between $\vec{A}$ and ψ .

Finally, though $\varphi : \pi_1(M) \rightarrow G$ is a group homomorphism, and $\tilde{M} \xrightarrow{f} M \times G$ given by $f(y) = (\pi(y), 1)$ is a canonical map, there is no bundle map between ξ_c and ξ : the pair of functions $(f \times \varphi, f)$ is *not* a principal bundle homomorphism.

In summary, the three bundles are related by the following diagram:

$$\begin{array}{ccccc} \pi_1(M) & \xrightarrow{\varphi} & G & & \mathbb{C}^m \\ \downarrow & & \downarrow & & \downarrow \\ \tilde{M} & \xrightarrow{f} & M \times G & \xrightarrow{\iota} & (M \times G) \times_G \mathbb{C}^m \\ \downarrow \pi & & \downarrow \pi_G & & \downarrow \pi_{\mathbb{C}^m} \uparrow \psi \uparrow \nabla_V^\omega \psi \\ M & = & M & = & M \end{array}$$

where ι is the canonical injection of the bundle ξ into its associated bundle i.e. $\iota(p) = [p, 0]$.

Acknowledgments

This work was partially supported by the grant PAPIIT-UNAM IN103505. M. S. thanks for hospitality to the Instituto de Astronomía y Física del Espacio (UBA-CONICET, Argentina), and the University of Valencia, Spain, where part of this work was done.

References

1. Aharonov, Y. and Bohm, D.(1959). Significance of Electromagnetic Potentials in Quantum Theory, *Physical Review*, **115**, 485-491.
2. Chambers, R. G. (1960). Shift of an Electron Interference Pattern by Enclosed Magnetic Flux, *Physical Review Letters*, **5**, 3-5.
3. Daniel, M. and Viallet, C. M. (1980). The Geometrical Setting of Gauge Theories of the Yang-Mills Type, *Reviews of Modern Physics*, **52**, pp. 175-197.
4. Aguilar, M. A. and Socolovsky, M. (2002). Aharonov-Bohm Effect, Flat Connections, and Green's Theorem, *International Journal of Theoretical Physics*, **41**, 839-860. (For a review see, e.g. ref. 5.)

5. Socolovsky, M. (2006). Aharonov-Bohm Effect, in *Encyclopedia of Mathematical Physics*, Elsevier, Amsterdam, pp. 191-198.
6. Wu, T. T. and Yang, C. N. (1975). Concept of non Integrable Phase Factors and Global Formulation of Gauge Fields, *Physical Review D*, **12**, 3845-3857.
7. Sundrum, R. and Tassie, L. J. (1986). Non-Abelian Aharonov-Bohm Effects, Feynman Paths, and Topology, *Journal of Mathematical Physics*, **27**, 1566-1570.
8. Botelho, L. C. L. and de Mello, J. C. (1987). A Non-Abelian Aharonov-Bohm Effect in the Framework of Feynman Pseudoclassical Path Integrals, *Journal of Physics A: Math. Gen.*, **20**, 2217-2219.
9. Harris, E. G. (1996). The gravitational Aharonov-Bohm effect with photons, *American Journal of Physics*, **64**, 378-383.
10. Zeilinger, A., Horne, M. A. and Shull, C. G. (1983). Search for unorthodox phenomena by neutron interference experiments, *Proceedings International Symposium of Foundations of Quantum Mechanics*, Tokyo, pp. 289-293.
11. Ho, Vu B. and Morgan, M. J. (1994). An experiment to test the gravitational Aharonov-Bohm effect, *Australian Journal of Physics*, **47**, 245-252.
12. Bezerra, V. B. (1987). Gravitational analogue of the Aharonov-Bohm effect in four and three dimensions, *Physical Review D*, **35**, 2031-2033.
13. Wisnivesky, D. and Aharonov, Y. (1967). Nonlocal effects in classical and quantum theories, *Annals of Physics*, **45**, 479-492.
14. Corichi, A. and Pierri, M. (1995). Gravity and Geometric Phases, *Physical Review D*, **51**, 5870-5875.
15. Greenberg, M. J. and Harper, J. R. (1981). *Algebraic Topology. A First Course*, Addison-Wesley, Redwood City, p. 126.
16. Nash, C. and Sen, S. (1983). *Topology and Geometry for Physicists*, Academic Press, London, p. 262.
17. Idem 16., pp. 147-148.
18. Naber, G. L. (2000). *Topology, Geometry, and Gauge Fields, Interactions*, Springer-Verlag, New York, pp. 193-197.
19. Massey, W. S. (1991). *A Basic Course in Algebraic Topology*, GTM 56, Springer, New York, p. 132.
20. Schulman, L. S. (1981). *Techniques and Applications of Path Integration*, Wiley, New York, pp. 205-207.
21. Kobayashi, S and Nomizu, K. (1963). *Foundations of Differential Geometry, Vol. 1*, Wiley, New York, p. 71.

e-mails:

rshuerfanob@unal.edu.co

alicia@nucleares.unam.mx

socolovs@nucleares.unam.mx